


# Xylazine Test Strips

These 4 steps could save your life. **1- CRUSH** **2- MEASURE** **3- TEST** **4- READ**

1

## Crush and mix.

Crush into fine powder and mix together.



1 scoop



## Measure and add water.


Add 1 microscop of your drug for every 5ml of water.

2


3

## Stir and test.

Stir well with microscop. Place wavy side of the strip into the water for 15 seconds.




Remove and place on a flat surface for 5 minutes.



## Read your results.

One red line is positive for xylazine.  
Two red lines is negative for xylazine.



4



If no red lines appear or one red line appears in the dotted area, the results are incomplete and should be repeated. **If your drugs contain cocaine, it will produce a false positive result.**

**Stop, think, and plan ahead.**

**Start low, go slow and let someone know.**



**CAUTION** - No test is 100% accurate. Your drugs may still contain fentanyl, xylazine and other drugs. **Consume drugs after testing at your own risk.**

# Testing Your Drugs

- Mixing different drugs unevenly can be dangerous.
- A sample of the drug test may not have the same contents as the rest of the supply. Dissolve your whole dose in water for testing.
- You can drink the water after testing or get the powder back by heating the water in a heat-safe container.
- Do not eat or consume test strips. Throw in garbage after use.

A negative result does not indicate drug safety. **Use drugs at your own risk.**

- Temperature may affect test accuracy. Use test strips at room temperature.
- Store test strips between 2°-30°C, and do not use if damaged, wet or expired.



## WARNING

No test is 100% accurate. Your drugs may still contain fentanyl, xylazine or other drugs.

## Overdose Protection

Street drugs may be mixed with other substances. Protect yourself from overdose.

- Avoiding mixing drugs.
- Start low, go slow — always try a small amount first.
- Always have naloxone kits on hand.
- Don't use alone.
- Call 9-1-1 in the event of an overdose.



**It is best to test everything you intend to ingest.**



## Naloxone

### Get a Naloxone Kit

Naloxone is a safe drug that can prevent and temporarily reverse the effects of an overdose caused by opioids.

Available **FREE** at the Health Unit, your local pharmacy or outreach organization.



### 5 Steps to Respond to an Overdose

1. Shout and shake
2. Call 9-1-1
3. Give naloxone
4. Perform rescue breathing and/or chest compressions
5. Check - Is it working?



Learn more about the Xylazine Test Strips:

[www.btnx.com/Product?id=2019](http://www.btnx.com/Product?id=2019)