


INSTRUCTION SHEET

Fentanyl Test Strips





IN CASE OF EMERGENCY

Testing Your Drugs

- Mixing different drugs unevenly can be dangerous.
- A sample of the drug test may not have the same contents as the rest of the supply. Dissolve your whole dose in water for testing.
- You can drink the water after testing or get the powder back by heating the water in a heat-safe container.
- Do not eat or consume test strips. Throw in garbage after use.

A negative result does not indicate drug safety. **Use drugs at your own risk.**

- Temperature may affect test accuracy. Use test strips at room temperature.
- Store test strips between 2°-30°C, and do not use if damaged, wet or expired.



WARNING

No test is 100% accurate. Your drugs may still contain fentanyl, xylazine or other drugs.

Overdose Protection

Street drugs may be mixed with other substances. Protect yourself from overdose.

- Avoiding mixing drugs.
- Start low, go slow always try a small amount first.
- Always have naloxone kits on hand.
- Don't use alone.
- Call 9-1-1 in the event of an overdose.





Get a Naloxone Kit

Naloxone is a safe drug that can prevent and temporarily reverse the effects of an overdose caused by opioids.

Available **FREE** at the Health Unit, your local pharmacy or outreach organization.



5 Steps to Respond to an Overdose

- 1. Shout and shake
- 2. Call 9-1-1
- 3. Give naloxone
- 4. Perform rescue breathing and/or chest compressions
- 5. Check Is it working?



Learn more about the Fentanyl Test Strips:

www.dancesafe.org/drug-checking

IN CASE OF EMERGENCY

CALL 9-1-1